

Asian Journal of Case Reports in Surgery

Volume 7, Issue 2, Page 528-531, 2024; Article no.AJCRS.126811

Abdominal Situs Inversus with Jejuno-Ileal Atresia and Polysplenia: A Case Report

Basel Maher ^a, Prince Raj ^{a*}, Raghu Shankar ^a and Martin Corbally ^a

^a Department of General Surgery, King Hamad University Hospital, Bahrain.

Authors' contributions

This work was carried out in collaboration among all authors. Authors PR and RS conceptualized and designed the study. Author MC played a role as the main surgeon of the participant. Authors BM, PR, RS and MC searched for Literature review, the consolidation of references and wrote the manuscript. Authors RS and PR conducted patient review, medical follow-up and performed the data collection. All authors read and approved the final manuscript.

Article Information

Open Peer Review History: This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/126811

Case Report

Received: 10/09/2024 Accepted: 14/11/2024 Published: 20/11/2024

ABSTRACT

The presence of an abnormally positioned organ is known as situs ambiguous. Polysplenia is defined as the presence of more than one spleen. This is a case of a baby girl antenatal diagnosed to have bowel dilatation and at birth presented with gross abdominal distention and bile-stained aspirate. Derotation of the distal apple peel ileal deformity and alignment of opposing mesentery and end-to-end jejuno-ileal anastomosis was done. Total small bowel length of 37 cms was preserved. Post-operative period was uneventful and she was gradually weaned from total parenteral nutrition to full feeds by day 17. An important aspect to consider is that derotation for volvulus is contrary to the normal derotation of malrotation / volvulus and should be performed in a

Cite as: Maher, Basel, Prince Raj, Raghu Shankar, and Martin Corbally. 2024. "Abdominal Situs Inversus With Jejuno-Ileal Atresia and Polysplenia: A Case Report". Asian Journal of Case Reports in Surgery 7 (2):528-31. https://journalajcrs.com/index.php/AJCRS/article/view/579.

^{*}Corresponding author: E-mail: drprp1610@gmail.com;

clockwise direction not anticlockwise. Other gastrointestinal anomalies associated with situs inversus are duodenal atresia, annular pancreas, biliary atresia, preduodenal portal vein, diaphragmatic hernia, lung cyst, genitourinary anomalies, ear, eye, and vertebral defects. Surgical principles remain same as for classic cases, though if there is any midgut volvulus derotation has to be performed in clockwise direction as volvulus in situs inversus will occur in anti-clockwise direction.

Keywords: Situs inversus; polysplenia; jejuno-ileal atrea.

1. INTRODUCTION

Situs inversus viscerum, is described as reversal of the internal body organs which can be complete (totalis) where both the thoracic and abdominal organs are reversed or might be partial (partialis) where either the thoracic or abdominal organs are reversed [1]. The presence of an abnormally positioned organ is known as situs ambiguous. Polysplenia is defined as the presence of more than one spleen. It is associated with situs inversus in about 20% of the cases . Jejuno-ileal atresia is a common cause of intestinal obstruction in neonates and its incidence is about 1 in 5000 live births. Having these three conditions in association is extremely rare.

2. CASE STUDY

This is a case of a baby girl antenatal diagnosed to have-bowel dilatation and at birth presented with gross abdominal distention and bile-stained aspirate. Abdominal radiograph showed dilated bowel loops (Fig 1(a)), and contrast enema revealed microcolon. At laparotomy multiple intestinal atresia (Fig 1(b)) with Type 3b jejunoileal atresia (apple peel appearance), situs Inversus abdominis and polysplenia (in right upper guadrant) were noted. Resection of 10cm of dilated jejunum just proximal to atresia was done along with resection of two sites of ileal atresia. Derotation of the distal apple peel ileal deformity and alignment of opposing mesentery and end-to-end ieiuno-ileal anastomosis was done. Total small bowel length of 37 cms was preserved. Proximal 10 cm of jejunum and distal 27 cm of ileum was preserved. Trans anastomotic feeding tube was not placed. Postoperative period was uneventful and she was gradually weaned from total parenteral nutrition to full feeds by day 17. An interrupted Inferior vena cava (IVC) and ridge in the descending aorta with a tiny inlet ventricular septal defect was noted on cardiac evalutaion. Further, the findings were confirmed with a CT Angiogram which showed focal mild narrowing of the aorta just distal to the left subclavian artery with mild post stenotic dilatation; there was an absent hepatic segment of the IVC with the hepatic veins draining into the subclavian vein. The baby was discharged on day 27 of life on full oral feeds.



Fig. 1 (a): Abdominal X-ray done pre-operatively with dilated bowel loops with the tip of orogastric tube in right hypochondrium; (b): Intra-operative findings showing multiple atresia

3. DISCUSSION

Situs inversus is rare with a prevalence of I in 10,000 of the general population. The genetics of situs inversus are complex and multifactorial and largely unknown though , maternal diabetes, retinoic acid exposure, conjoined twinning, cocaine abuse, and consanguinity are all implicated. The embryology has not been well established however, the rotational anomaly can cause a vascular accident due to midgut volvulus during intrauterine period which could be the cause of multiple atresia. Situs inversus is generally asymptomatic and is diagnosed incidentally but when this is associated with atresias or midgut volvulus, presentation is guite soon after birth in the neonatal period [2]. An important aspect to consider is that derotation for volvulus is contrary to the normal derotation of malrotation / volvulus and should be performed in a clockwise direction not anticlockwise. Other gastrointestinal anomalies associated with situs inversus are duodenal atresia, annular pancreas, preduodenal biliarv atresia. portal vein. diaphragmatic hernia, lung cyst, genitourinary anomalies, ear, eye, and vertebral defects [3].

Ileal and jejunal atresias are usually described collectively as jejuno-ileal atresia and is are causes common of neonatal intestinal obstruction. Grosfeld described four types of intestinal atresia; Type IIIb and type IV as in our case, is with absence of the distal superior mesenteric artery; and distal small bowel coiling (like an apple peel) [4]. Bowel loss is present with multiple atresias, which was the case as noted intra-operatively in our patient. Poor outcomes are associated with prematurity, low birth weight, and associated anomalies in type IIIb jejuno-ileal atresia.

Sporadic cases have described the presence of polysplenia, situs inversus abdominus and multiple intestinal atresia and are rarely reported in the English literature [5,6]. Apart from the case presented here, only a single case report could be retrieved highlighting the rare association of these three entities [7]. Though in the case reported by Chinya et al situs inversus was diagnosed pre-operatively and they sited their abdominal incision accordingly, but most commonly these associations are encountered intra-operatively when operating for intestinal atresia or midgut volvulus. Though our preoperative abdominal x-ray was suspicious of situs inversus abdominis it was missed (tip of orogastric tube was in right hypochondrium). We

don't feel that knowing this pre-operatively would have changed our management plan or for that matter the siting of incision would be of great importance as for exploration we need to have adequate exposure, but definitely if known prior to surgery intra-op surprises could be minimized. Challenge in these situation is not the situs abdominis or the multiple spenenculi but it's the multiple atresia with Type 3b association. There is always a risk of short bowel with Type 3b, and when this is associated with multiple atresia we need to be extra careful in how much bowel to resect. The length of dilated proximal jejunum and the length of remaining distal ileum should be adequate enough to prevent any short gut syndrome.

4. CONCLUSION

Situs inversus in association with polysplenia and multiple JIA is an extremely rare association-Preoperative diagnosis of situs inversus is often missed in spite of some tell-tale sign. Plain radiographs should be carefully looked for in cases with malrotation or jejunoileal atresias. Surgical principles remain same as for classic cases, though if there is any midgut volvulus derotation has to be performed in clockwise direction as volvulus in situs inversus will occur in anti-clockwise direction.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT

Informed consent for publication was taken from the parents

ETHICAL APPROVAL

It's not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ruben GD, Templeton JM, Ziegler MM: Situs inversus: the complex inducing

neonatal intestinal obstruction. J Pedia Surg. 1983;18:751-6.

- Abdur-Rahman LO, Adeniran JO, Taiwo JO. Concurrent dextrogastria, reverse midgut rotation and intestinal atresia in a neonate. J Indian Assoc Paediatr Surg. 2007;12:228–230.
- Sharma S, Rashid KA, Dube R, Malik G K, Tandon R K. Congenital duodenal obstruction with situs inversus totalis: Report of a rare association and discussion. J Indian Assoc Pediatr Surg. 2008;13:77-8
- Dalla Vecchia, Laura K., et al. "Intestinal atresia and stenosis: a 25year experience with 277 cases. " Archives of Surgery. 133.5 (1998): 490-497.
- Nawaz A, Matta H, Hamchou M, Jacobez A, Trad O, Al Salem AH. Situs inversus abdominus in association with congenital duodenal obstruction: a report of two cases and review of the literature. Pediatr Surg Int. 2005;21:589–92.
- 6. Rasool, Farooq. Bilal and Mirza. "Polysplenia syndrome associated with situs inversus abdominus and type I jejunal atresia. " APSP Journal of Case Reports. 2011;2(2):18.
- Chinya, Abhishek et al. Situs inversus abdominalis, polysplenia, complex jejunal atresia and malrotation in a neonate: A rare association. International Journal Of Surgery Case Reports. 2019;56:93-95.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/126811